References:

Aggarwal BB, Anand P, Sundaram C, Jhurani S, Kunnumakkara AB (2008). Curcumin and cancer: An ‘‘old-age” disease with an ‘‘age-old” solution. Cancer Letters. 267:133–164.

Alonson MJ, Lorenzo-Lamosa ML, Remunan-Lopez C, Vila-Jato JL (1998). Design of microencapsulated chitosan microspheres for colonic drug delivery. J. Control. Release. 52: 109–118.

Anderson DMW, Weiping W (1990). The composition of some Sesbania gum exudates. Biochemical Systematics and Ecology . 18(1):43-44.

Ansari MJ, Ahmad S, Kholi K, Ali J, Khar RK (2005). Stability-indicating HPTLC determination of curcumin in bulk drug and pharmaceutical formulations. Journal of Pharmaceutical and Biomedical Analysis. 39:132-138.

Ash AJ (1990). The effect of supplementation with leaves from the leguminous trees Sesbania grandiflora, Albizia chinensis and Gliricidia sepium on the intake and digestibility of guinea grass hay by goats. Animal Feed Science and Technology. 28(3-4): 225-232.

Bhaskar RPR, Krishnaiah YSR, Satyanarayana V, Karthikeyan RS (2002). Studies on the development of oral colon targeted drug delivery systems for metronidazole in the treatment of amoebiasis. Int. J. Pharm. 236 (1-2): 43–55.

Brondsted H, Andersen C, Hovgaard L (1998). Crosslinked dextran – a new capsule material for colon targeting of drug. J. Control. Release. 53: 7–13.

Bussemer T, Otto I, Bodmeier R (2001). Pulsatile drug delivery systems: Critical Review. Ther Drug Carrier Syst. 18(5): 433-458.

Cheng AL, Hsu CH, Lin JK, Hsu MM, Ho YF, Shen TS (2001). Phase I clinical trial of Curcumin, a chemopreventive agent, in patients with high-risk or pre-malignant lesions, Anticancer Res. 21 (4B):2895–2900.

Dusel R, McGinity J, Harris M R, Vadino W A, Cooper J (1986). Sodium alginate, Handbook of Pharmaceutical Excipients, Published by American pharmaceutical Society: USA and The Pharmaceutical Society of Great Britain, London: 257-258.

Gayathri P, Satheshkumar PS, Prasad K, Nair S, Savithri HS, Murthy MRN (2006). Crystal structure of the serine protease domain of Sesbania mosaic virus polyprotein and mutational analysis of residues forming the S1-binding pocket. Virology. 346:440-451.

Giri B, Mukerji KG (2004). Mycorrhizal inoculant alleviates salt stress in Sesbania aegyptiaca and Sesbania grandiflora under field conditions: evidence for reduced sodium and improved magnesium uptake. 14(5):307-312.

Gothoskar AV, Joshi AM, Joshi NH (2004). "Pulsatile Drug Delivery Systems: A Review", Drug Delivery Technology. 4(5): 121-125.

Gurudu S, Fiocchi C, Katz J (2002). Inflammatory bowel disease. Best Practice and Research Clinical Gastroenterology. 16(1):77-90.

Haeberlin B, Empey L, Fedorak R, Nolen H, Friend D R (1993). In vivo studies in the evaluation of glucuronide prodrug for novel therapy of ulcerative colitis. Proceedings of the International Symposium on Controlled Release of Bioactive Materials. 20:174-175.

Hebden JM, Wilson CG, Spiller RC, Gilchrist PJ, Blackchaw E, Frier ME, Perkin AC (1999). Regional difference in quinine absorption from the undistributed human colon assessed using a timed release delivery system. Pharma. Res. 16: 1087-1092.

Heineman M, Otieno JO, Mengich EK, Amadalo BA (1997). Growth and yield of eight agroforestry tree species in line plantings in Western Kenya and their effect on maize yields and soil properties. Forest Ecology and Management. 91(1):103-135.

Hodson CA, Mitchell JR, Davies MC, Melia CD (1995). Structure and behavior in hydrophilic matrix dosage forms: The influence of pH on the sustained release performance and internal gel structure of sodium alginate matrices. J. Control. Release. 33(1): 143-152.

Hossain MA, Focken U, Becker K (2002). Nutritional evaluation of dhaincha (*Sesbania aculeata*) seeds as dietary protein source for tilapia *Oreochromis niloticus.* 33(9):653-662.

Hovgaard L, Brondsted H (1996). Current application of polysaccharides in colon targeting: Critical Review. Ther. Drug Carrier Syst. 13: 185-223.

Ireson C, Orr S, Jones DJL, Verschoyle R, Lim CK, Luo JL (2001). Characterization of metabolites of the chemopreventive agent curcumin in humans and rat hepatocytes and in

the rat in vivo, and evaluation of their ability to inhibit phorbol ester-induced prostaglandin E2 production. Cancer Res. 61:1058–1064.

Jing Lei, Ji-Heung Kim (2008). Preparation and Properties of Alginate/Polyaspartate Composite Hydrogels. Macromolecular Research. 16 (1): 45-50.

Kasture VS, Deshmukh VK, Chopde CT (2002). Anxiolytic and anticonvulsive activity of Sesbania grandiflora leaves in experimental animals. Phytother Res. 16(5): 455-460.

Kawamori T, Lubet R, Steele VE, Kelloff GJ, Kaskey RB, Rao CV (1999). Chemopreventive effect of curcumin, a naturally occurring anti-inflammatory agent, during the promotion/progression stages of colon cancer, Cancer Res. 59 (3): 597–601.

Kholi K, Ansari J, Reheman Z (2005). Curcumin: A natural antiinflammatory agent. Indian Journal of Pharmacology. 37: 141-147.

Kiyoung L, Kun N, Yueim K (1999). Polysaccharides as a drug coating polymer. Polym. Prep. 40: 359–360.

Krishnaiah YSR, RamaPrasad YV, Satyanarayana S (1998). In vitro evaluation of guar gum as a carrier for colon-specific drug delivery. J. Control. Release. 51(2-3): 281–287.

Lakshminarayana R, Raju M, Krishnakantha TP, Baskaran V (2005). Determination of major carotenoids in a few Indian leafy vegetables by high-performance liquid chromatography. J. Agric. Food Chem. 53(8):2838-2842.

Lal B, Khanna S (1994). Selection of salt-tolerant Rhizobium isolates of Acacia nilotica . World Journal of Microbiology and Biotechnology . 10 (6): 637-639.

Lee S, Lim CB, Lee S, Pai CM (1999). Colon selective drug delivery. Composion. WO99/01115.

LiuXinga, ChenDaweia, XieLipingb, ZhangRongqingb (2003). Oral colon-specific drug delivery for bee venom peptide: development of a coated calcium alginate gel beads-entrapped liposome. J. Controlled Release. 93: 293–300.

Maheshwari RK, Singh AK, Gaddipati J, Srimal RC (2006). Multiple biological activities of curcumin: A short review. Life Sciences. 78: 2081–2087.

Mandal AS, Biswas N, Karim KM, Guha A, Chatterjee S, Behera M, Kuotsu K(2010). Drug delivery system based on chronobiology—A review. J. Control. Release. 147(3): 314–325.

Maurice O, Iwunze (2004).Binding and distribution characteristics of curcumin solubilized in CTAB micelle. Journal Molecular Liquids. 111(1-3):161-165.

Milojevic S, Newton J M, Cummings J H, Gibson G R, Botham R L, Ring S G, Stockham M, Allwood M C (1995). Amylose, the new perspective in oral drug delivery to the human large intestine. S.T.P. Pharm. Sci. 51: 47–53.

Milojevic S, Newton J M, Cummings J H, Gibson G R, Botham R L, Ring S G, Stockham M, Allwood M C (1996a). Amylose as a coating for drug delivery to the colon: preparation and in vitro evaluation using 5-aminosalicylic acid pellets. J. Control. Release. 38(1): 75–84.

Mishra B, Jayanth P, sankar C (2003). Development of chitosan-alginate microcapsules for colon specific delivery of metronidazole. Indian Drugs. 40(12):695-700.

Momin RK, Kadam VB (2011). Determination of Lipid and Alkaloid Content in Some Medical Plant. 3(3):18–21.

Motterlini R, Foresti R, Bassi R, Green CJ (2000). Curcumin, an antioxidant and anti-inflammatory agent, induces heme oxygenase-1 and protects endothelial cellsagainst oxidative stress. Free Radical Biology & Medicine. 28 (8): 1303–1312.

Murthy MRN , Bhuvaneswari M, Subramanya HS, Gopinath K, Savithri HS (1997). Structure of Sesbania mosaic virus at 3 Å resolution. Biophysical Chemistry. 68 (1-3): 33-42.

Negi PS, Jayaprakash GK, Jagan MRL, Sakariah KK (1999). Antibacerial activity of turmeric oil; A byproduct from curcumin manufactuer. J.Agric.Food Chem. 47: 4297-300.

Onsoyen E, Karsa DR (1995). Hydration induced swelling of alginate based matrix tablets at GI tract pH conditions. Excipients and delivery systems for pharmaceutical formulations. The royal society of chemistry, Cambridge.108-122.

Paradkar A, Anshuman A, Ambike, Bhimrao K, Jadhav, Mahadik KR (2004). Characterization of curcumin–PVP solid dispersion obtained by spray drying. International Journal of Pharmaceutics. 271(1-2): 281-286.

Pari L, Uma A (2003). Protective effect of Sesbania grandiflora against erythromycin estolate-induced hepatotoxicity. Therapie. 58(5):439-443.

Pertuit D, Moulari B, Betz T, Nadaradjane A, Neumann D, Ismaïli L, Refouvelet B, Pellequer Y, Lamprecht A (2007). 5-amino salicylic acid bound nanoparticles for the therapy of inflammatory bowel disease. J. of Control. Release. 123 (2007) 211–218.

Pillay V, Fassihi R (1999a). In vitro release modulation from crosslinked pellets for site-specific drug delivery to the gastrointestinal tract. I. Comparison of pH responsive drug release and associated kinetics. J. Controlled Release. 59(2): 229-242.

Pillay V, Fassihi R (1999b). In vitro release modulation from crosslinked pellets for site-specific drug delivery to the gastrointestinal tract. I. Physicochemical characterization of calcium-alginate, calcium pectinate and calcium-alginate- pectinate pelletsa. J. Controlled Release. 59(2): 243-256.

Pollard MA, Fischer P, Windhab EJ (2011). Characterization of galactomannans derived from legume endosperms of genus Sesbania (Faboideae). 84 (1): 550-559.

Radai R, Rubinstein A (1995). In vitro and in vivo analysis of colon specificity of calcium pectinate formulations. Eur. J. Pharm. Biopharm. 41: 291–295.

Ramsewak RS, DeWitt DL, Nair MG (2000). Cytotoxicity, antioxidant and anti-inflammatory activities of curcumins I–III from Curcuma longa-Phytomedicine. 7(4): 303–308.

Rubinstein A, Nakar D, Sintov A (1992a). Chondroitin sulphate: a potential biodegradable carrier for colon-specific drug delivery. Int. J. Pharm. 1992a; 84(2): 141–150.

Rubinstein A, Nakar D, Sintov A (1992b). Colonic drug delivery: enhanced release of indomethacin from crosslinked chondritin matrix in rat cecal content. Pharm. Res. 9(2): 276–278.

Rubinstein A, Radai R, Ezra M, Pathak S, Rokem JS (1993). In vitro evaluation of calcium pectinate: a potential colon-specific drug delivery carrier. Pharm. Res. 10(2): 258–263.

Ruby AJ, Kuttan G, Dinesh BK, Rajasekharan KN, Kuttan R (1995).Antitumor and antioxidant activity of natural Curcuminoids. Cancer Lett. 94: 79–83.

Satyanaryana S, Krishnaiah YSR, Rama Prasad YV (1999). Studies of guar gum compression-coated 5-aminosalicylic acid tablets for colon-specific drug delivery. Drug Dev. Ind. Pharm. 25: 651–657.

Sharma RA, Ireson CR, Verschoyle RD (2001). Effects of dietary curcumin on glutathione S-transferase and malondialdehyde DNA adducts in rat liver and colon mucosa: relationship with drug levels. Clin. Cancer Res. 7:1452–1458.

Shinha VR, Kumaria R (2004). Polysaccharide matrix for microbially triggered drug delivery to the colon. Drug Dev. Ind. Pharm. 30(2):143-150.

Shivakumar HG, Pramod kumar TM, Kashppa GD (2003). "Pulsatile drug delivery system", Indian J. Pham. Educ. 37(3): 125.

Shun YL, Ayres JW (1992). Calcium Alginate beads as core carriers of 5-Aminosalicylic acid. Pharm. Res. 9: 714–790.

Singh KI, Singh J, Sharma D, Sharma A (2012). Colon Specific Drug Delivery System: Review on Novel Approaches. Int. J. Pharm. Sci. & Res. 3(3): 637-647.

Sinha VR, Kumaria R (2001). Polysaccharide in colon specific drug delivery. Int. J. Pharm. 224:19-38.

Skaugrund O, Karsa DR, Stephenson RA (1995). Drug delivery system with alginate and chitosan. Excipients and delivery systems for pharmaceutical formulations. The royal society of chemistry, Cambridge. 95-107.

Sowbhagya HB, Smitha S, Sampathu SR, Krishnamurthy N, Bhattacharya S (2005). Journal of food engineering. 67:367-371.

Sugavara S, Imai T, Otagiri M (1994). The controlled release of prednisolone using alginate gel. Pharm. Res. 2: 272-277.

Tominaga S, Takaizawa T, Yamada M (1998). Colon drug delivery system. J. Pol. Appl. 10: 324- 642.

Tønnesen HH, Másson M, Loftsson T (2002). Studies of curcumin and curcuminoids. XXVII. Cyclodextrin complexation: solubility, chemical and photochemical stability. Int. J. Pharm. 244(1-2):127-135.

Turkoglu M, Ugurlu T (2002). In vitro evaluation of pectin–HPMC compression coated 5-aminosalicylic acid tablets for colonic delivery. Eur. J. Pharm. Biopharm. 53(1): 65–73.

Vervoort L, Rambant P, Van den Mooter G, Augustijns P, Kinget R (1998b). Inulin hydrogels. II. In vitro degradation study. Int. J. Pharm. 172(1-2):137–145.

Vervoort L, Van den Mooter G, Augustijins P, Busson R, Toippet S, Kinget R (1997). Inulin hydrogels as carriers for colonic drug targeting: I. Synthesis and characterization of methacrylated inulin and hydrogel formation. Pharm. Res. 14: 1730–1737.

Vervoort L, Van den Mooter G, Augustijns P, Kinget R (1998a). Inulin hydrogels. I. Dynamic and equilibrium swelling properties. Int. J. Pharm. 172(1-2): 127–135.

Vervoort L, Vinckier I, Moldenaers P, Van den Mooter G, Ahgustigns P, Kinget R (1999). Inulin hydrogels as carriers for colonic drug targeting. Rheological characterization of the hydrogel formation and the hydrogel network. J. Pharm. Sci. 88: 209.

Vidyasagar S, Ramakrishna BS (2002). Effects of butyrate on active sodium and chloride transport in rat and rabbit distal colon. Journal of Physiology. 539 (1): 163–173.

Vyas SP, Khar RK (2002). Controlled Drug delivery concepts and advances. Vallabh Prakashan, 1st ed.:234-255.

Wang YJ, Pan MH, Chang AL, Hsieh CY, Lin JK (1997). Stability of curcumin in buffer solutions and characterization of its degradation products. J. Pharm. Biomed. Anal. 15(12):1867-1876.

Wilding IR, Davis SS, Wilding EA (1993). Gastrointestinal transit of a matrix tablet formulation: comparison of canine and human data. Int. J. Pharm. 94 (1-3): 235-238.

Yang L, Chu JS, Fix JA (2002). Colon-specific drug delivery: New approaches and in vitro/in vivo evaluation. Int. J. Pharm. 235 (1-2):1–15.

Youan B, Harris M (2004). The biological requirement of a given disease therapy J. Control. Rel. 98:337-353.